2003 Health and Preventive Practices Status Report

HEALTH AND SENIOR SERVICES.CHRONIC DISEEASE PREVENTION AN

D HEALTH PROMOTION

Clay County**

from Missouri County-level Study

	County Prevalence (%)*	R.C. Metro Region Prevalence (%)*	State Prevalence (%)*
General			
Health Status (Fair or Poor)	13.0	15.2	16.9
No Health Coverage	6.8 L	10.3	12.3
Behavioral Risk Factors			
Physical Inactivity	20.6	25.9	23.0
Current Smoker	22.1	23.4	26.5
Overweight (BMI 25-29.9)	31.5	35.6	35.3
Obese (BMI ≥ 30)	32.0	27.4	23.4
D _{isease and Conditions}			
High Blood Pressure	27.4	27.9	28.5
High Cholesterol (35 years or older)	37.2	43.3 H	37.3
Asthma	11.3	12.7	10.7
Diabetes	4.7	6.8	7.2
Activity Limitation	14.2	18.7	17.8
Cancer Screening			
Women			
Never had a Mammogram	17.5	13.0	13.4
No Mammogram or Clinical Breast Exam in Last Year:			
40-49 years of age	58.1	48.0	49.4
50-64 years of age	17.3	40.4	35.9
65 and older	38.3	43.7	47.4
Never had a Pap Test:			
18-69 years of age	2.4	3.3	5.1
70 and older	2.6	3.1	6.4
No Pap Test in Last Year (18-29)	12.7	23.5	25.8
No Pap Test in Last 3 Years (30-69)	12.3	9.1	13.8
Men and Women 50 Years or Older			
Blood Stool Test:			
Never had a Blood Stool Test	51.9	46.2 L	57.0
No Blood Stool Test in Last Year	68.4	72.0	77.5
Sigmoidoscopy/Colonoscopy			
Never had a Sigmoidoscopy or Colonoscopy	40.8	40.8	49.2

County

K.C. Metro

State



*Prevalence rates are age adjusted to the 2000 U.S. standard population

indicates that the prevalence rate is not significantly above or below state prevalence

49.2

48.8

57.0

- indicates that the prevalence rate is significantly above the state prevalence
- indicates that the prevalence rate is significantly below the state prevalence

Unless otherwise noted respondents are 18 years of age or older

No Sigmoidoscopy or Colonoscopy in Past Five Years

^{**} The best indicators for Clay county were obtained by combining data from Clay and Platte counties.